

# The Diode Laser as an Electrosurgery Replacement

by Glenn A. van As, BSc, DMD

## Introduction

In 2008, Dr. Gordon Christensen wrote an article in *JADA* comparing the soft tissue cutting ability of diode lasers versus that of electrosurgery (radiosurgery) units.<sup>1</sup> In that article, he compared these two technologies against each other, and cited advantages and disadvantages of each alternative. In choosing between the two technologies at that time, he made several points:

1. Although, there was considerable overlap in their uses, and both technologies were effective, Christensen found there were a few potential uses that did not overlap. These included the use of diode lasers around metal (amalgam and gold) as well as implants, that lasers did not harm dental hard tissues (bone) or soft tissues (pulp), that the clinician could use the laser with less anesthetic, and finally that lasers were antimicrobial (antibacterial).
2. The use of the laser, especially the diode laser, was increasing in dentistry, and that lasers attract patients because of their recognized and accepted role within the field of medicine (LASIK eye surgery).
3. Electrosurgery units were “far less expensive than the least expensive diode lasers” and he questioned whether “the advantages of the diode laser significant enough to compensate for the additional cost.”

Now, less than two years later, a tremendous surge in diode laser interest has occurred. This is in due in no small part to the

drop in the price range of diode lasers from around the \$10,000-\$12,000 in 2008 for the least expensive diode semiconductor lasers to a price range of \$2,500 (see AMD Lasers’ Picasso Lite). This dramatic price drop of more than 75 percent in the price of these units, has allowed diode lasers to become less expensive than some bipolar electrosurgery units and comparable, but still more expensive than many monopolar electrosurgery units. These monopolar electrosurgery units can be purchased for \$1,000 or less, but the question as to whether the added benefits of diode lasers cited by Christensen are now enough to make them a soft-tissue alternative to these units for the average clinician. In this article I will review these questions again, and suggest that for many dental practices a simple diode laser might replace their electrosurgery unit as the methodology of choice for soft-tissue laser surgery.

## Advantages of Diode Lasers Over Electrosurgery

As Christensen mentioned in his article, the term “electrosurgery” is not as well known or as accepted by many patients as the term “lasers.” Having said this, the acceptance of this technology in both medicine and dentistry, as a viable method of soft-tissue alteration has made electrosurgery an accepted alternative to the scalpel. There are two basic types of electrosurgical units:



**Monopolar**, in which a single electrode exists and the current travels from the unit down a single wire to the surgical site. The patient is grounded with a pad placed behind the patients back (a part of the procedure that many patients may question). Heat is produced when the electrode contacts the tissue, and due to pain that is produced anesthetic must be used.

**Bipolar**, in which two electrodes that are in very close proximity exist on these units. Bipolar units are more expensive than diode lasers and the electrical current flows from one electrode to the other, thus eliminating the need for a grounding pad. Bipolar units, because of the two wires, create less of a precise cut than the monopolar or diode laser. Electrosurgical units will typically cut larger amounts of tissue with greater speed than diode lasers can, and this can be important if very large amounts of tissue must be removed. In many cases, with routine dentistry, soft-tissue ablation is of the minor to moderate amount in nature, and in this case speed is not an issue.

### Tissue Troughing with the Diode Laser

The diode laser has become a popular technology as an alternative for tissue management compared to the traditional methodology of placing a single or double retraction cord in the sulcus. Many CEREC users routinely use the diode laser to enhance the gingival trough when the margin is either equi-gin-

gival or subgingival, prior to powdering their prep for their digital impression. The diode laser can be used in almost all instances to produce gingival retraction as an alternative to cord with excellent results both in terms of gingival retraction and margin delineation for the laboratory. Diodes, like electrosurgical units, offer the clinician the ability to work in a bloodless field for the impressions because of the hemostasis that occurs during the procedure. Unlike electrosurgical units where recession can be an issue, as can postoperative pain, diode lasers offer the clinician the ability to precisely remove overhanging, inflamed tissue while creating a gingival trough that is not likely to cause damage to bone, cementum, or pulp tissue like electrosurgical units can.

A small learning curve exists, in knowing when and how to properly diode laser trough (*see clinical case 1 and table 1*). Once the learning curve is conquered most clinicians will almost completely eliminate cord from their practice. In critically aesthetic areas where thin tissue genotypes exist, or if the patient is changing the color of the tooth significantly from the existing stump shade, then care with diode troughing must be taken.

In the author's experience, with the introduction of adequate levels of magnification (Loupes 4.0X or greater or an operating microscope) and the careful use of lower powers on the diode laser (for example 0.6-0.9 watts of power in Continuous Wave), the diode laser tissue management can be done with confidence in not having gingival recession occur post-operatively, and often can be done with only topical anesthetics. This is particularly enticing in situations where the tooth has had previous endodontic therapy and tissue troughing can be completed with the use of stronger topical anesthetics (Cetycaine, Tricaine Blue, TAC 20, EMLA) and lower settings with the laser. Use of the electrosurgery unit mandates the need for chemical anesthetic (injections of local anesthetic) in order to complete the tissue troughing. In addition, there is research that suggests that the lateral thermal damage done with lasers is significantly lower than that with electrosurgery.

**Table 1: Seven-step Clinical Procedure for Laser Crown Troughing**

1. Initial gross reduction and margin placement equi-gingival
2. Diode laser troughing: suggested settings 0.6-1.1 w CW (less in anterior)
3. Final margin placement subgingivally as needed for aesthetics
4. Hydrogen Peroxide or wet cotton pellet to remove tissue tags
5. Lateral distention of tissue if needed (Expasyl, Traxodent)
6. Rinse and take PVS impression
7. Provisional fabrication – careful to make sure no overhangs

*continued on page 58*

### Clinical Case 1: Comparing Diode Laser Troughing to Retraction Cord




*Fig. 1: Preop of cracked premolars needing full coverage restorations.  
 Fig. 2: Diode laser being used to “laser trough” around first premolar.  
 Fig. 3: Retraction cord being placed on second premolar of case.  
 Fig. 4: Occlusal view of two teeth treated with alternative technologies.  
 Fig. 5: Low magnification view of impression.  
 Fig. 6: High magnification view showing differences in impression of gingival sulcus with cord (left) and laser (right) but acceptable results with both methods.  
 Fig. 7: Occlusal view of healed tissue after provisionals removed after two weeks.  
 Fig. 8: LAVA crowns in place on both teeth – occlusal view immediately post-op.  
 Fig. 9: LAVA crowns in place on both teeth – labial view immediately post-op.*

The safety of lasers for gingival retraction procedures has been documented in the literature by Gherlone, et. al.<sup>2</sup> who found that lasers (diode and NdYAG) when compared to the conventional techniques of double cord or electro-surgery

yielded less gingival bleeding, and also less gingival recession. Their interesting conclusion was that although both techniques are satisfactory that the laser techniques were in fact “less traumatic to the periodontal tissues.”

### Diode Laser Usage Around Metals

The diode laser has the added benefit of being able to be used with less concern over damage to hard or soft dental tissues, or damage to dental prosthesis that can occur with the less expensive monopolar electro-surgery units.<sup>3-6</sup> The diode laser can be used safely around gold crowns and amalgam restorations without fear of pulpal damage (Figs. 10-14) and lasers can be safely used around implants with minimal fear of introducing iatrogenic damage to the implant or bone, and in new research there are suggestions that the diode lasers when used at lower levels of power (Low Level Laser Therapy) may in fact improve the early healing of tissues and in also be used in cases of peri-implantitis (Figs. 15-20).<sup>7-14</sup>



*Fig. 10: Large buccal amalgam on UR 2nd Molar.*

*Fig. 11: Amalgam partially removed – gingivectomy needed to remove rest of amalgam.*

*Fig. 12: Laser gingivectomy completed without bleeding or fear of sparking.*

*Fig. 13: Rest of amalgam restorative material removed.*

*Fig. 14: Restoration completed on labial.*

*Fig. 15: Flipper partial over top of implant healing cap for right upper central incisor.*

*Fig. 16: Healing cap in place after Flipper partial removed.*

*Fig. 17: Topical on tissue to allow for gingivectomy around implant on mesial to provide for provisional crown to be seated properly.*

*Fig. 18: Immediate postoperative view after laser gingivectomy.*

*Fig. 19: Provisional crown placed.*

*Fig. 20: Note healing of tissue prior to final PFM crown placement (Three months healing of provisional crown).*

*Fig. 21: Final crown in place.*

## Use of Anesthetic

One of the key benefits of the diode laser over electro-surgery units is the reduced need for local anesthetics. Some minor recontouring for aesthetics and for orthodontics can be accomplished with only topical anesthetic. The literature shows that lasers can safely be used for both cosmetic and orthodontics<sup>15-18</sup> and there are an increasing number of clinicians buying lasers for this very fact alone. As I am fond of saying, “Dentists are fine with needles and drills... it’s the patients who are seeking alternatives.” Electrosurgery units and scalpels almost always require the usage of local anes-

*continued on page 60*

**AFFORDABLE**  
*Dental*  
**MARKETING**

- DEMOGRAPHIC ANALYSES
- LOGOS & PROMO ITEMS
- STATIONERY & INTERNALS
- RECALL & REFERRAL PIECES
- WEBSITES & HOSTING
- TURN-KEY DIRECT MAIL
- PRINT ADS OF ALL SIZES
- BANNERS & PROMO BOARDS
- PRACTICE BROCHURES

**0%-INTEREST FINANCING** Call **1-888-575-2233**  
**www.PracticeCafe.com**

FREE FACTS, circle 2 on card


thetic. Figures 22-30 show cases of cosmetic and orthodontics done with no local anesthetic.

**Cosmetic “Smile Lifts”**

Lasers have been shown to be effective in cosmetic dentistry cases.<sup>19-22</sup> The lure of the laser is to help with “gummy smile” cases where an excessive or asymmetrical amount of tissue appears in a smile. All too often, we as dentists focus on the “white” parts of the smile and fail to observe gingival asymmetries (pink part of the smile) which, if corrected,

could significantly improve the overall aesthetic outcome of the case.

Conventional periodontal surgery consists of full or partial thickness flaps, and osseous surgery to remove bone, followed by sutures and 12-16 weeks of healing. There is nothing wrong with viewing the overall architecture of the underlying biology of bone, roots and soft tissue, however there are times when more minimally invasive techniques may be used with equally impressive results but with perhaps much less healing time.



*Fig. 22: Pre-op view of canines requiring exposure of clinical crown to place orthodontic bracket.*

*Fig. 23: Higher magnification of upper right canine.*

*Fig. 24: Topical gel (compounding pharmacy) applied to labial tissue.*

*Fig. 25: Diode exposure of clinical crown with 1.2-1.4 w pulsed setting.*

*Fig. 26: Final view of brackets in place on both upper canines.*

*Fig. 27: Pre-op view of pigmented lesion on attached tissue noticed by patient.*

*Fig. 28: Higher magnification view of lesion.*


*Fig. 29: Immediate postoperative appearance after laser ablation at 1.2w pulsed.*

*Fig. 30: Ten day healing photo of tissue.*

*Fig. 31: High magnification view of healing showing complete disappearance of lesion.*

One of the causes of “short tooth syndrome” where a lack of the clinical crown is displayed when smiling is altered passive eruption. In these cases, the osseous level subgingival has receded apical to the CEJ. The gingival margins in Active Passive Eruption (APE) has not moved coronal enough to the level of the CEJ. This often leaves large amounts of Keratinized tissue which can be removed via a diode laser gingivectomy.

In cases where a “gummy smile” exists, the clinical crowns are short, but the bone is not apical to the CEJ, (altered active eruption is one cause), then diode laser gingivectomies will not lead to a stable clinical result. Osseous surgery either in a full flap scenario or at times with closed flap laser techniques using both diode and erbium lasers, can provide tremendous results for the patient with shortened healing times (Figures 32-37).<sup>23-25</sup>



*Fig. 32: Preoperative smile shows “small tooth syndrome.”*

*Fig. 33: Shows existing 3/4 porcelain crowns on maxillary incisors.*

*Fig. 34: Shows Er:YAG closed flap crown lengthening (smile lift) and gingivectomies immediately post-op.*

*Fig. 35: Shows diode laser tissue management four weeks afterwards prior to impressions.*

*Fig. 36: Postoperative result shows longer clinical crowns and optimum tissue health.*

*Fig. 37: Shows closeup of final result.*

continued on page 62

## Anti-bacterial Capabilities of Lasers

Many articles in the literature have demonstrated the tremendous ability of all lasers with respect to bacterial and even fungal reduction.<sup>26-33</sup> This feature alone makes lasers effective and desirable in many areas in the oral cavity where the risk of postoperative infection may be reduced with lasers. Particular interest is now occurring with the role of lasers in endodontic, periodontic and peri-implantitis cases where the need to reduce bacterial loads without such a great deal of reliance on antibiotics might be exciting. Although more research is needed on how the bactericidal capabilities of the diode laser might be beneficial in these areas, there is no debating that all lasers can help healing through decreasing the risk of infection through laser light alone (Figs. 38-42). In addition, growing research has demonstrated that the risk of high bacterial loads in periodontal pockets and in particular in endodontics situations may be reduced by lasers. These newer articles have implications for improving traditional methodologies locally where used, and in helping to reduce the potential greater systemic health risks generally. The role of lasers continues to be researched today, but present research has shown that diode lasers can be used safely within root canals with minimal fear of developing iatrogenic complications when conservative settings are used.<sup>34-38</sup>

**Negotiating a Lease?**

Don't let the landlord take advantage of you.




**“You are a champion in your field!”**

- Dr. Dan Myers  
Aphareta, G A

To level the playing field, simply call toll-free or visit [georgevail.com/dt/](http://georgevail.com/dt/)

**GEORGEVAIL**  
DENTAL OFFICE LEASE NEGOTIATIONS  
**800-340-2701**





*Fig. 38: Four canal upper second molar prior to obturation.  
 Fig. 39: 400 micron diode fiber in DB canal for bacterial reduction.  
 Fig. 40: Diode laser for bacterial reduction in canal.  
 Fig. 41: Final view of obturated four canals in second molar.  
 Fig. 42: Preoperative and postoperative radiographs of above case*

**Treatment of Oral Lesions**

One of the advantages of a diode laser is the ability to treat oral lesions including: Recurrent Aphthous Ulcers (RAU), Venous Lake lesions of the lips and Herpetic Lesions (Figs. 43-49). Research has shown that lasers can be safely used to treat these lesions,<sup>39-41</sup> and in addition it is possible that if caught early during the prodromal stage that the lesions can be aborted or significantly reduced in terms of length of time they are present.<sup>42</sup> In addition, it has been the author's experience that, once treated with the laser, the lesions are often less likely to reappear in the same area.

Venous Lake lesions of the lower lip have been traditionally one of the more difficult lesions to treat. There is a growing body of evidence to show that a diode laser can often without anesthetic completely eliminate these purplish lesions which occur frequently on the lips in one single treatment, often with only topical anesthetic.<sup>14,15, 43, 44</sup>

*Fig. 43: Pre-op appearance of venous lake on lower right lip.  
 Fig. 44: High mag preoperative appearance.  
 Fig. 45: Diode laser coagulating venous lake (note uninitiated tip).  
 Fig. 46: Immediate postoperative appearance of lesion.  
 Fig. 47: High mag appearance of venous lake immediately postoperative.  
 Fig. 48: One-week healing of lip-lesion considerably smaller.  
 Fig. 49: Two-week healing of lip (lesion is now fully disappeared).*



**Conclusion**

In the last two years, diode lasers have become a staple of many dental practices for their cost effective solution to many clinical problems that are seen daily in private practice. The laser as an "electrosurgery replacement" has become a reality with

these newer units, which provide numerous advantages to everyday dentistry.

The advantages of diode laser tissue troughing as a replacement in many instances for cord, in being safely used around metals (implants, amalgam, gold, orthodontic brackets), hard and soft tissue, cannot be overlooked. In many instances small amounts of tissue can be removed with only topical anesthetics (Cetycaine, EMLA, Tricaine Blue, and TAC 20) and diode lasers are great soft-tissue lasers for many orthodontic and cosmetic procedures as well. If one adds in the antibacterial capabilities of these lasers and their ability to be used in many soft-tissue surgeries including frenectomies, fibroma removals, and the treatment of oral lesions, then it can be seen that perhaps the soft tissue laser is on its way to being an important and essential part of not only every dental practice, but perhaps in the not-too-distant future, an integral part of every dental operator. ■

### Author's Bio

**Dr. Glenn van As** graduated from the University of British Columbia in 1987 and is internationally known for digital documentation of laser procedures captured with the operating microscope. He achieved Advanced Proficiency from the Academy of Laser Dentistry and received the 2006 Leon Goldman Award for clinical excellence. He is also a founder and past president of the Academy of Microscope Enhanced Dentistry.



### References

1. Christensen GJ. Soft-tissue cutting with laser versus electrosurgery. *J Am Dent Assoc.* 2008 Jul;139(7):981-4.
2. Gherlone EF, Maionara C, Grassi RF, Ciacaglini R, Cattoni F. The use of 980-nm Diode and 1064-nm Nd:YAG Laser for Gingival Retraction in Fixed Prosthesis. *J Oral Laser Applications.* 2004; 4:183-190.
3. Robertson PB, Luscher B, Spangberg LS, Levy BM. Pulpal and periodontal effects of electrosurgery involving cervical metallic restorations. *Oral Surg Oral Med Oral Pathol.* 1978 Nov;46(5):702-20.
4. Yoshino T, Aoki A, Oda S, Takasaki AA, Mizutani K, Sasaki KM, Kinoshita A, Watanabe H, Ishikawa I, Izumi Y. Long-term histologic analysis of bone tissue alteration and healing following Er:YAG laser irradiation compared to electrosurgery. *J Periodontol.* 2009 Jan;80(1): 82-92.
5. Skaria AM. Electrocoagulation and hazardous damage to a dental prosthesis. *J Am Acad Dermatol.* 2006;54(3):543-4.
6. Wilcox CW, Wilverding TM, Watson P, Morris JT. Use of electrosurgery and lasers in the presence of dental implants. *Int J Oral Maxillofac Implants* 2001;16(4): 578-82.
7. Yeh S, Jain K, Andreeana S. Using a diode laser to uncover dental implants in second-stage surgery. *Gen Dent.* 2005 Nov-Dec;53(6):414-7.
8. Hauser-Gerspach I, Stubinger S, Meyer J. Bactericidal effects of different laser systems on bacteria adhered to dental implant surfaces; and in vitro study comparing zirconia with titanium. *Clin Oral Implants Res.* 2010 Jan 13.
9. Stubinger S, Homann F, Etter C, Miskiewicz M, Wieland M, Sader R. Effect of Er:YAG, CO(2) and diode laser irradiation on surface properties of zirconia endosseous dental implants. *Lasers Surg Med.* 2008 Mar;40(3):223-8.
10. Giangormus M, Akyol U. The effect of gallium-aluminum-arsenide 808-nm low-level laser therapy on healing of skin incisions made using a diode laser. *Photomed Laser Surg.* 2009 Dec; 27(6): 895-9.
11. AboElsaud NS, Soory M, Gadalla LM, Ragab LI, Dunne S, Zalata KR, Louca C. Effect of soft laser and bioactive glass on bone regeneration in the treatment of infra-bony defects (a clinical study). *Lasers Med Sci.* 2009 May; 24(3):287-95. Epub 2008 Jun 26.
12. Matsumoto MA, Ferino RV, Monteleone GV, Ribeiro DA. Low-level laser therapy modulates cyclo-oxygenase-2 expression during bone repair in rats. *Laser Med Sci.* 2009 Mar;24(2):195-201. Epub 2008 Feb 29.
13. Castro GL, Gallas M, Nunez IR, Borrado JL, Alvarez JC, Varela GL. Scanning electron microscopic analysis of diode laser-treated titanium implant surfaces. *Photomed Laser Surg.* 2007 Apr;25(2):124-8.
14. Dorthbudak O, Haas R, Mallath-Pokorny G. Biostimulation of bone marrow cells with a diode soft laser. *Clin Oral Implants Res.* 2000 Dec;11(6):540-5.

continued on page 64



**WOULD GETTING YOUR DENTAL PRACTICE ON THE FIRST PAGE OF GOOGLE®, YAHOO® AND BING® PUT A SMILE ON YOUR FACE?**

Whether you need a customized website or you already have one that you want to be seen on the first page of search results, **call us today!**

**ONE DENTAL OFFICE PER ZIP CODE AVAILABLE!**

FOR A LIMITED TIME WE ARE OFFERING CUSTOMIZED WEBSITES FOR ONLY **\$499** Ask for promo code: **TOWNIES**

- Custom Designed Websites
- Edit Your Site for FREE
- Patient Educational Videos
- Top Search Engine Results (Limit one office per zip code)
- Professional Email Addresses

endorsed by



**www.DentistIdentity.com • 800.303.6029**

15. Azevedo LH, Galletta VC, de Paula Eduardo C, Migliari DA. Venous Lake of the Lips Treated Using Photocoagulation with High-Intensity Diode Laser. *Photomed Laser Surg.* 2010 Apr;28(2):263-5.
16. Genovese WJ, dos Santos MT, Faloppa F, de Souza Merli LA. The use of surgical diode laser in oral hemangioma: a case report. *Photomed Laser Surg.* 2010 Feb;28(1):147-51.
17. Fornaini C, Rocca JP, Bertrand MF, Merigo E, Nammour S, Vescovi P. Nd:YAG and diode laser in the surgical management of soft tissues related to orthodontic treatment. *Photomed Laser Surg.* 2007 Oct;25(5):381-92.
18. Genovese MD, Olivi G. Use of laser technology in orthodontics: hard and soft tissue laser treatments. *Eur J Paediatr Dent.* 2010 Mar;11(1):44-8.
19. Pang P. Lasers in cosmetic dentistry. *Gen Dent.* 2008 Nov-Dec;56(7):663-70.
20. Magid KS, Straus RA. Laser use for esthetic soft tissue modification. *Dent Clin North Am.* 2007 Apr;51(2):525-45.
21. Meeks T. Creating beautiful smile symmetry: tissue considerations. *Dent Today.* 2009 Oct;28(10):98,100-1.
22. Adams TC, Pank PK. Lasers in aesthetic dentistry. *Dent Clin North Am.* 2004 Oct;48(4):833-69, vi.
23. Lowe RA. Minimally invasive dentistry combined with laser gingival plastic surgery: maximize your aesthetic results. *Dent Today.* 2008 Aug;27(8):102, 104-5.
24. Eshom DS. The Er,Cr:YSGG laser periodontal surgery. *Prac Proced Aesthet Dent.* 2008 Aug;20(7):433-5.
25. Hornbrook DS. In the spotlight: three of the hottest topics in dentistry! Interview by Damon Adams. *Dent Today.* 2009 Sep;28(9):94, 96, 98-9.
26. Kuvvetli SS, Sandalli N, Topcuoglu N, Kulekci G. Antibacterial efficacy of diode and Er:YAG laser irradiation in experimentally contaminated primary root canals. *J Clin Pediatr Dent.* 2009 Fall; 34(1): 43-8.
27. Cobb CM, Low SB, Coluzzi DJ. Lasers and the Treatment of Chronic Periodontitis. *Dent Clin N Am.* 2010; 54:35-53.
28. Aoki A, Mizutani K, Takasaki AA, et al. Current status of clinical laser applications in periodontal therapy. *Gen Dent* 2008;56(7):674-87.
29. de Souza EB, Cai S, Simionato MR, Lage-Marques JL. High-power diode laser in the disinfection in the depth of the root canal dentin. *Oral Surg Oral Med Oral Pathol Radiol Endod.* 2008 Jul;106(1):e68-72.
30. Sennhenn-Kirchner S, Schwarz p, Schliephake H, Konieschke F, Brunner E, Borg-von Zepelin M. Decontamination efficacy of erbium:yttrium-aluminium-garnet and diode laser light on oral *Candida albicans* isolates of a 5-day in vitro biofilm model. *Laser Med Sci.* 2009 May;24(3):313-20. Epub 2008 May 6.
31. Schoop U, Kluger W, Devisbegovic S, Goharkhay K, Wernisch J, Georgopoulos A, Sperr W, Moritz A. Innovative wavelengths in endodontic treatment. *Lasers Surg Med* 2006 Jul;38(6):624-30.
32. Gutknecht N, Franzen R, Schippers M, Lampert F. Bactericidal effect of a 980-nm diode laser in the root canal wall dentin of bovine teeth. *J Clin Laser Med Surg.* 2004 Feb;22(1):9-13.
33. Schoop U, Kluger W, Moritz A, Nedjelik N, Georgopoulos A, Sperr W. Bactericidal effect of different laser systems in the deep layers of dentin. *Laser Surg Med.* 2004;35(2): 111-6.
34. Hmud R, Kabler WA, George R, Walsh LJ. Cavitational effects in aqueous endodontic irrigants generated by near-near infrared lasers. *J Endo.* 2010 Feb;36(2): 275-8. Epub 2009 Dec.4.
35. Alfredo E, Marchesan MA, Sousa-Neto MD, Brugnara-Junior A, Silva-Sousa YT. Temperature variation at the external root surface during 980-nm diode laser irradiation in the root canal. *J Dent.* 2008 Jul;36(7):529-34. Epub 2008 May 6.
36. Gutnecht N, Franzen R, Meister J, Vanweersch L, Mir M. Temperature evolution on human teeth root surface after diode laser assisted endodontic treatment. *Laser Med Sci.* 2005 Sep;20(2):99-103. Epub 2005 Jul 9.
37. Theodoro LH, Haypek P, Bachmann L, Garcia VG, Sampaio JE, Zezell DM, Eduardo Cde P. Effect of ER:YAG and Diode laser irradiation on the root surface: morphological and thermal analysis. *J Periodontol.* 2003 Jun;74(6): 838-43.
38. da Costa Ribeiro A, Nogueira GE, Antoniazzi JH, Moritz A, Zezell DM. Effects of diode laser (810 nm) irradiation on root canal walls: thermographic and morphological studies. *J Endod.* 2007 Mar;33(3):252-5. Epub 2006 Dec 13.
39. Kravitz ND, Kusnoto B. Soft-tissue lasers in orthodontics: an overview. *Am J Orthod Dentofacial Orthop.* 2008 Apr;133(4 Suppl):S110-4.
40. Tezel A, Kara C, Balkaya V, Orbak R. An Evaluation of Different Treatments for Recurrent Aphthous Stomatitis and Patient Perceptions: Nd:YAG Laser versus Medication. *Photomed Laser Surg.* 2009 Feb;27(1):101-6.
41. Colvard M, Kuo P. Managing aphthous ulcers: laser treatment applied. *J Am Dent Assoc.* 1991 Jun;122(6):51-3.
42. Kotlow L. Lasers and pediatric dental care. *Gen Dent.* 2008 Nov-Dec;56(7): 618-27.
43. Rice JH. Removal of venous lake using a diode laser (810nm). *Wavelengths* 2004;12(1):20-1.
44. Kotlow LA. Elimination of a venous lake on the lower vermillion of the lower lip via 810-nm diode laser. *J Laser Dent* 2007;15(1):20-22.

## DOCTORS CHOICE GOLD EXCHANGE

Step 1  
Request a Scrap  
for Cash-Pak!

Step 2  
Send us your Scrap\*,  
Coins or Jewelry!

Step 3  
WE PAY YOU!  
IT'S THAT EASY!

**Doctors Choice Gold Exchange was originated by a "Dentist" who was frustrated at the low prices paid to consumers, primarily dentists, who turned in their scrap gold only to receive a small percentage of its actual worth from numerous companies promising highest prices.**

**Credibility • Integrity • Ethics**  
Visit: [Doctorschoicgoldexchange.com](http://Doctorschoicgoldexchange.com)  
To receive "Top Dollar" for your precious metal scrap!

**HERE'S WHAT YOU DO:**

*Gather your unwanted Gold, Precious and Semi-Precious material, whether it is Dental Scrap\*, Jewelry or Coins that you want to convert to CASH! Fill out the online information form on the left of any page to get your insured FREE SCRAP for CASH-PAK.*

**HERE'S WHAT WE DO:**

*We receive your material and immediately verify the weight and purity of all precious and semi-precious material. Once this is done, we base our value of your material on the P.M. London Gold Fix at the time of receipt of your material. We put more profit in your pocket!*

**HERE'S WHAT WE DON'T DO:**

*We don't send this to a middleman or agent thus diluting the profits, profits intended for YOU! That is why Doctors Choice Gold Exchange can pay the high returns to YOU the CONSUMER!*

**1-877-279-2601**

**Doctorschoicgoldexchange.com**

©2009 Doctors Choice Gold Exchange, L.L.C. All Rights Reserved